



**NH Division of Public Health Services
Immunizations Required for Childcare Attendance**

Vaccine	Recommended Schedule	Comment
DTaP (Diphtheria, Tetanus, a cellular pertussis)	2 months, 4 months, 6 months, booster 15-18 months, booster 4-6 years.	Age appropriately required for childcare attendance per routine childhood vaccination schedule.
Polio eIPV (enhanced inactivated polio vaccine) OPV (oral polio vaccine)	2 months, 4 months, 6-18 months, booster 4-6 years	Age appropriately required for childcare attendance per routine childhood vaccination schedule.
Haemophilus influenzae Type b (HbOC) ActHIB (4 doses) (PRP-T) PedvaxHIB (PedvaxHIB 3 doses)	2 months, 4 months, 6 months, booster 12-15 months	Age appropriately required for childcare attendance. If the child starts the series late or is behind, fewer doses may be required. One dose at 15 months or older is adequate. Not required after 59 months of age.
MMR (measles, mumps, rubella)	1 dose administered on or after 12 months of age. 2 nd dose recommended 4-6 years of age.	Age appropriately required for childcare attendance per routine childhood vaccination schedule. 2 nd dose must be at least 1 month after dose one.
Varivax (chickenpox)	1 dose administered on or after 12 months of age. 2 nd dose recommended 4-6 years of age.	Age appropriately required for childcare attendance per routine childhood vaccination schedule. 2 nd dose must be at least 3 months after dose one. (For children ages 12 months to 12 years).
Hep B (hepatitis-B vaccine)	Birth – 2 months, 2-4 months, booster 6-18 months.	Age appropriately required for childcare attendance.

**NH Division of Public Health Services
Immunizations Recommended for Childcare Attendance**

Vaccine	Recommended Schedule	Comment
Hep A (Hepatitis A vaccine)	12 months with a booster 6 months later.	Age appropriate recommended for childcare attendance per routine childhood vaccination schedule.
Influenza vaccine	6-months-18 years, one dose annually.	If it's the first time receiving vaccine, two doses are required separated by one month. See child's primary health care provider for further guidance.
PCV-7 (7 valent pneumococcal conjugated vaccine)	2 months, 4 months, 6 months, booster 12-15 months.	Age appropriately recommended for childcare attendance, per routine childhood vaccination schedule. If the child starts the series late or is behind, fewer doses may be required. Recommended 23-59 months in certain high- risk groups.
Rotavirus vaccine (Two different vaccine products)	Rotarix (RV1) 2 months, 4 months. Rotateq (RV5) 2 months, 4 months, 6 months.	Age appropriate recommended for childcare attendance per routine childhood vaccination schedule.

*1 month is equal to 28 days

** Derived from the *Harmonized Immunization Schedule* approved by the American Academy of Pediatrics, the American Academy of Family Physicians, and the Advisory Committee on Immunization Practices.

If you have any questions about a child's compliancy, please call the child's primary care provider or the New Hampshire Immunization Program (1-800-852-3345 Ext. 4482)